

Hamiltonian formulation of systems with linear velocities within Riemann-Liouville fractional derivatives

February 4, 2008

Sami I. Muslih ¹

Department of Physics, Al-Azhar University, Gaza, Palestine
and
International Center for Theoretical Physics(ICTP),
Trieste, Italy

Dumitru Baleanu ²

Department of Mathematics and Computer Sciences, Faculty of Arts and
Sciences, Çankaya University- 06530, Ankara, Turkey

Abstract

The link between the treatments of constrained systems with fractional derivatives by using both Hamiltonian and Lagrangian formulations is studied. It is shown that both treatments for systems with linear velocities are equivalent.

PACS: 11.10.Ef

Mathematics Subject Classification: 26A33, 70Hxx, 70H50

Key words: fractional derivative, Hamiltonian system, Non-conservative systems.

¹E-mail: smuslih@ictp.trieste.it

²On leave of absence from Institute of Space Sciences, P.O.BOX, MG-23, R 76900, Magurele-Bucharest, Romania, E-mails: dumitru@cankaya.edu.tr, baleanu@venus.nipne.ro

1 Introduction

The generalization of the concept of derivative and integral to a non-integer order α has been subjected to several approaches and some various alternative definitions of fractional derivatives appeared [1-6]. In the last few years fractional calculus was applied successfully in various areas , e.g. chemistry , biology, modelling and identification, electronics and wave propagation. Fractional calculus , has played an important role in engineering, science, and pure and applied mathematics [7-9]. Fractional derivatives were applied in recent studies of scaling phenomena [10-12]. Classical mechanics is one of the fields when fractional calculus found many applications [13-19]. Riewe has used the fractional calculus to obtain a formalism which can be applied for both conservative and non conservative systems [13, 14]. Although many laws of nature can be obtained using certain functionals and the theory of calculus of variations, not all laws can be obtained by this manner. As it is known, almost all systems contain internal damping, yet traditional energy based approach cannot be used to obtain equations describing the behavior of a non conservative system [13,14]. Using the fractional calculus one can obtain the Lagrangian and the Hamiltonian equations of motion for the nonconservative systems .

The understanding of constrained dynamics [20], both at the classical and quantum level, has been a subject of long standing theoretical interest, which has seen important contributions ever since Dirac's quantization of the electromagnetic field. The path integral approach and the canonical one are two main approaches of quantization.

Recently, an extension of the simplest fractional problem and the fractional variational problem of Lagrange was obtained [17,18]. Even more recently, this approach was extended to Lagrangians with linear in velocities [21,22] , which represents a typical example of second-class constrained systems in Dirac's classification [20]. These Lagrangians are important because their Euler-Lagrangian equations become systems of first order differential equations in contrast with second order corresponding to the regular ones. In addition, these systems may possess gauge symmetries and gauge ambiguities.

From these reasons it is interesting to study the fractional Hamiltonian of constrained systems.

The aim of this paper is to obtain the fractional Hamiltonian equations of motion for Lagrangians with linear velocities.

The plan of our paper is as follows:

In Sect. 2 some basic tools of fractional derivatives as well as Riewe's approach of the fractional Lagrangian and Hamiltonians are presented. In Sect. 3 the Euler-Lagrange equations were obtained using the Agrawal's approach and the fractional formulation of systems with constraints is introduced. In Sect 4 the fractional Hamiltonian of the systems possessing linear velocities is analyzed. Conclusion is presented in section 5.

2 Fractional Lagrangian and Hamiltonian formulations

In this section we briefly present the definition of the left and right derivatives together with Riewe's formulation of Lagrangian and Hamiltonian dynamics. The left Riemann-Liouville fractional derivative is defined as

$${}_a\mathbf{D}_t^\alpha f(t) = \frac{1}{\Gamma(n-\alpha)} \left(\frac{d}{dt}\right)^n \int_a^t (t-\tau)^{n-\alpha-1} f(\tau) d\tau, \quad (1)$$

and the right Riemann-Liouville fractional derivative has the form

$${}_t\mathbf{D}_b^\alpha f(t) = \frac{1}{\Gamma(n-\alpha)} \left(-\frac{d}{dt}\right)^n \int_t^b (\tau-t)^{n-\alpha-1} f(\tau) d\tau, \quad (2)$$

where the order α fulfills $n-1 \leq \alpha < n$ and Γ represents the Euler's gamma function. If α is an integer, these derivatives are defined in the usual sense, i.e.,

$${}_a\mathbf{D}_t^\alpha f(t) = \left(\frac{d}{dt}\right)^\alpha, \quad {}_t\mathbf{D}_b^\alpha f(t) = \left(-\frac{d}{dt}\right)^\alpha, \quad \alpha = 1, 2, \dots \quad (3)$$

Now we shall briefly review Riewe's formulation of fractional generalization of Lagrangian and Hamiltonian equations of motion [13,14]. The starting point is the action function of the form

$$S = \int_a^b L(\{q_n^r, Q_{n'}^r\}, t) dt. \quad (4)$$

Here the generalized coordinates are defined as

$$q_n^r := ({}_a\mathbf{D}_t^\alpha)^n x_r(t), \quad Q_{n'}^r := ({}_t\mathbf{D}_b^\alpha)^{n'} x_r(t), \quad (5)$$

and $r = 1, 2, \dots, R$ represents the number of fundamental coordinates, $n = 0, \dots, N$, the sequential order of the derivatives defining the generalized coordinates q , and $n' = 1, \dots, N'$ the sequential order of the derivatives in definition of the coordinates Q . A necessary condition for S to posses an extremum for given functions $x_r(t)$ is that $x_r(t)$ fulfill the Euler-Lagrange equations [13,14]

$$\frac{\partial L}{\partial q_0^r} + \sum_{n=1}^N ({}_t\mathbf{D}_b^\alpha)^n \frac{\partial L}{\partial q_n^r} + \sum_{n'=1}^{N'} ({}_a\mathbf{D}_t^\alpha)^{n'} \frac{\partial L}{\partial Q_{n'}^r} = 0. \quad (6)$$

Using the references [13,14], the generalized momenta have the following form

$$\begin{aligned} p_n^r &= \sum_{k=n+1}^N ({}_t\mathbf{D}_b^\alpha)^{k-n-1} \frac{\partial L}{\partial q_k^r}, \\ \pi_{n'}^r &= \sum_{k=n'+1}^{N'} ({}_a\mathbf{D}_t^\alpha)^{k-n'-1} \frac{\partial L}{\partial Q_k^r}. \end{aligned} \quad (7)$$

Thus, the canonical Hamiltonian is given by

$$H = \sum_{r=1}^R \sum_{n=0}^{N-1} p_n^r q_{n+1}^r + \sum_{r=1}^R \sum_{n'=0}^{N'-1} \pi_{n'}^r Q_{n'+1}^r - L. \quad (8)$$

The Hamilton's equations of motion are as follows

$$\frac{\partial H}{\partial q_N^r} = 0, \quad \frac{\partial H}{\partial Q_{N'}^r} = 0. \quad (9)$$

For $n = 1, \dots, N$, $n' = 1, \dots, N'$ we have the following equations of motion

$$\frac{\partial H}{\partial q_n^r} = {}_t\mathbf{D}_b^\alpha p_n^r, \quad \frac{\partial H}{\partial Q_{n'}^r} = {}_a\mathbf{D}_t^\alpha \pi_{n'}^r, \quad (10)$$

$$\frac{\partial H}{\partial q_0^r} = -\frac{\partial L}{\partial q_0^r} = {}_t\mathbf{D}_b^\alpha p_0^r + {}_a\mathbf{D}_t^\alpha \pi_0^r. \quad (11)$$

The remaining equations are given by

$$\frac{\partial H}{\partial p_n^r} = q_{n+1}^r = {}_a\mathbf{D}_t^\alpha q_n^r, \quad \frac{\partial H}{\partial \pi_{n'}^r} = Q_{n'+1}^r = {}_t\mathbf{D}_b^\alpha Q_{n'}^r, \quad (12)$$

$$\frac{\partial H}{\partial t} = -\frac{\partial L}{\partial t}, \quad (13)$$

where, $n = 0, \dots, N$, $n' = 1, \dots, N'$.

3 Fractional Euler-Lagrange equations

Recently Agrawal, have obtained the Euler-Lagrange equations for fractional variational problems [17]. In the following we like to present briefly his approach.

Consider the action function

$$S[q_0^1, \dots, q_0^R] = \int_a^b L(\{q_n^r, Q_{n'}^r\}, t) dt, \quad (14)$$

subject to the independent constraints

$$\Phi_m(t, q_0^1, \dots, q_0^R, q_n^r, Q_{n'}^r) = 0, \quad m < R, \quad (15)$$

where the generalized coordinates are defined as

$$q_n^r := ({}_a\mathbf{D}_t^\alpha)^n x_r(t), \quad Q_{n'}^r := ({}_t\mathbf{D}_b^\beta)^{n'} x_r(t). \quad (16)$$

Then, the necessary condition for the curves q_0^1, \dots, q_0^R with the boundary conditions $q_0^r(a) = q_0^{ra}, \quad q_0^r(b) = q_0^{rb}, \quad r = 1, 2, \dots, R$, to be an extremal of the functional given by equation (14) is that the functions q_0^r satisfy the following Euler-Lagrange equations [17]:

$$\frac{\partial \bar{L}}{\partial q_0^r} + \sum_{n=1}^N ({}_t\mathbf{D}_b^\alpha)^n \frac{\partial \bar{L}}{\partial q_n^r} + \sum_{n'=1}^{N'} ({}_a\mathbf{D}_t^\beta)^{n'} \frac{\partial \bar{L}}{\partial Q_{n'}^r} = 0, \quad (17)$$

where \bar{L} has the form [17]

$$\bar{L}(\{q_n^r, Q_{n'}^r\}, t, \lambda_m(t)) = L(\{q_n^r, Q_{n'}^r\}, t) + \lambda_m(t) \Phi_m(t, q_0^1, \dots, q_0^R, q_n^r, Q_{n'}^r). \quad (18)$$

Here the multiple $\lambda_m(t) \in R^m$ are continuous on $[a, b]$.

3.1 Fractional Hamiltonian formulation of Agrawal's approach

In order to obtain the Hamilton's equations for the the fractional variational problems, we re-define the left and the right canonical momenta as :

$$\begin{aligned} p_n^r &= \sum_{k=n+1}^N ({}_t\mathbf{D}_b^\alpha)^{k-n-1} \frac{\partial \bar{L}}{\partial q_k^r}, \\ \pi_{n'}^r &= \sum_{k=n'+1}^{N'} ({}_a\mathbf{D}_t^\beta)^{k-n'-1} \frac{\partial \bar{L}}{\partial Q_k^r}. \end{aligned} \quad (19)$$

Using (19), the canonical Hamiltonian becomes

$$\bar{H} = \sum_{r=1}^R \sum_{n=0}^{N-1} p_n^r q_{n+1}^r + \sum_{r=1}^R \sum_{n'=0}^{N'-1} \pi_{n'}^r Q_{n'+1}^r - \bar{L}. \quad (20)$$

Then, the modified canonical equations of motion are obtained as

$$\{q_n^r, \bar{H}\} = {}_t\mathbf{D}_b^\alpha p_n^r, \quad \{Q_{n'}^r, \bar{H}\} = {}_a\mathbf{D}_t^\alpha \pi_{n'}^r, \quad (21)$$

$$\{q_0^r, \bar{H}\} = {}_t\mathbf{D}_b^\alpha p_0^r + {}_a\mathbf{D}_t^\alpha \pi_0^r, \quad (22)$$

where, $n = 1, \dots, N$, $n' = 1, \dots, N'$.

The other set of equations of motion are given by

$$\{p_n^r, \bar{H}\} = q_{n+1}^r = {}_a\mathbf{D}_t^\alpha q_n^r, \quad \{\pi_{n'}^r, \bar{H}\} = Q_{n'+1}^r = {}_t\mathbf{D}_b^\alpha Q_{n'}^r, \quad (23)$$

$$\frac{\partial \bar{H}}{\partial t} = -\frac{\partial \bar{L}}{\partial t}. \quad (24)$$

Here, $n = 0, \dots, N$, $n' = 1, \dots, N'$ and the commutator $\{, \}$ is the Poisson's bracket defined as

$$\{A, B\}_{q_n^r, p_n^r, Q_{n'}^r, \pi_{n'}^r} = \frac{\partial A}{\partial q_n^r} \frac{\partial B}{\partial p_n^r} - \frac{\partial B}{\partial q_n^r} \frac{\partial A}{\partial p_n^r} + \frac{\partial A}{\partial Q_{n'}^r} \frac{\partial B}{\partial \pi_{n'}^r} - \frac{\partial B}{\partial Q_{n'}^r} \frac{\partial A}{\partial \pi_{n'}^r}, \quad (25)$$

where, $n = 0, \dots, N$, $n' = 1, \dots, N'$.

4 Equivalence of fractional Hamiltonian and Lagrangian formulations for systems with linear velocities

Recently, for $0 < \alpha \leq 1$, the Lagrangians with linear velocities were investigated in [21]. For example, the Euler-Lagrange equations of the following Lagrangian

$$L' = a_j (q^i) {}_a\mathbf{D}_t^\alpha q^j - V(q^i), \quad (26)$$

were obtained as [21]

$$\frac{\partial a_j(q^i)}{\partial q^k} {}_a\mathbf{D}_t^\alpha q^j + {}_t\mathbf{D}_b^\alpha a_k(q^i) - \frac{\partial V(q^i)}{\partial q^k} = 0. \quad (27)$$

Now we would like to obtain the Hamiltonian equations of motion for the same model. Let us define

$$x_j^n = ({}_a\mathbf{D}_t^\alpha)^n q_j, \quad n = 0, 1, \dots, N-1, \quad j = 1, 2, \dots, R. \quad (28)$$

The generalized momenta are given by,

$$p_j^0 = \frac{\partial L}{\partial x_j^0} = a_j(x_i^0), \quad , p_j^1 = \frac{\partial L}{\partial x_j^1} = 0. \quad (29)$$

The canonical Hamiltonian reads as

$$H = (p_j^0 - a_j(x_i^0)) x_j^1 + V(x_i^0). \quad (30)$$

The Hamiltonian equations of motion are calculated as

$$\frac{\partial H}{\partial x_j^1} = p_j^0 - a_j(x_i^0) = ({}_t\mathbf{D}_b^\alpha) p_j^1 = 0. \quad (31)$$

In fact, this equation is the primary constraint in the Dirac's formalism. The other equations of motion are calculated as

$$\frac{\partial H}{\partial x_k^0} = -\frac{\partial a_j(x_i^0)}{\partial x_k^0} x_j^1 + \frac{\partial V(x_i^0)}{\partial x_k^0} = ({}_t\mathbf{D}_b^\alpha) p_k^0, \quad (32)$$

$$\frac{\partial H}{\partial p_k^0} = x_k^1 = ({}_a\mathbf{D}_t^\alpha) q_k \quad (33)$$

Making use of equation (31) we obtain

$$\frac{\partial a_j(q^i)}{\partial q^k} {}_a\mathbf{D}_t^\alpha q^j + {}_t\mathbf{D}_b^\alpha a_k(q^i) - \frac{\partial V(q^i)}{\partial q^k} = 0. \quad (34)$$

It is obvious the equivalence between equations (34) and (27). An interesting point to be specified here is that, primary constraints in the Dirac's formalism are present as equations of motion in our treatment, while the other Hamiltonian equations of motion are equivalent to the Lagrangian equations of motion as given in [21].

5 Conclusion

One of the main problems encountered in applying the fractional calculus to a given singular Lagrangian and Hamiltonian is the existence of multiple choices of the possible fractional generalizations . In addition, the solutions of the fractional Euler-Lagrange equations contain more information than the classical ones. In this paper, Hamiltonian equations have been obtained for systems with linear velocities, in the same manner as those obtained by using the formulation of Euler-lagrange equations for variational problems introduced by one of us [21]. We study the general model for systems with linear velocities and it was observed that the Hamiltonian and Lagrangian equations which are obtained by the two methods are in exact agreement.

Acknowledgments

S. M. would like to thank the Abdus Salam International Center for Theoretical Physics, Trieste, Italy, for support and hospitality during the preparation of this work. This work was done within the framework of the Associateship Scheme of the Abdus Salam ICTP.

D. B. would like to thank O. Agrawal and J. A. Tenreiro Machado for interesting discussions. This work is partially supported by the Scientific and Technical Research Council of Turkey.

References

- [1] K. S. Miller and B. Ross, *An Introductation to the Fractional Integrals and Derivatives-Theory and Applications*, John Willey and Sons, New York, (1993).
- [2] S. G. Samko, A. A. Kilbas and O. I. Marichev, *Fractional Integrals and Derivatives- Theory and Applications*, Gordon and Breach, (1993).
- [3] K. B. Olham and J. Spanier, *The fractional calculus*, Academic Press, New York, (1974).
- [4] R. Gorenflo and F. Mainardi, *Fractional calculus: Integral and Differential Equations of Fractional Orders, Fractals and Fractional Calculus in Continoum Mechanics*, Springer Verlag, Wien and New York (1997).

- [5] I. Podlubny, *Fractional Differential Equations*, Academic Press, New York, (1999).
- [6] R. Hilfer, *Applications of Fraction Calculus in Physics*, World Scientific Publishing Company, Singapore, New Jersey, London and Hong Kong, (2000).
- [7] J.A. Tenreiro Machado, *A probabilistic interpretation of the fractional order differentiation*. Fract. Calc. Appl. Anal. 8(1)(2003),73-80.
J.A. Tenreiro Machado, *Discrete-time fractional -order controllers*, Fract. Calc. Appl. Anal. 4(1) (2001) 47-68.
- [8] O.P. Agrawal, Analytical solution for stochastic response of a fraction-ally damped beam, *Transactions of the ASME Journal of Vibration and Acoustics*, (2004), to appear.
- [9] M. Klimek, *Fractional sequential mechanics-models with symmetric frac-tional derivatives*, Czech. J. Phys. **51**, (2001), 1348-1354; *Lagrangian and Hamiltonian fractional sequential mechanics*, **52** (2002), 1247-1253.
- [10] T. F. Nonnenmacher, *Fractional integral and differential equations for a class of Levy-type probability densities*, J. Phys. **A 23** (1990), L697S-L700S.
- [11] R. Metzler and K. Joseph, *Boundary value problems for fractional dif-fusion equations*, Physica A. **278** (2000), 107-125 .
- [12] G. M. Zaslavsky, *Chaos, fractional kinetics, and anomalous transport* , Phys. Rep. **371** (2002), 461-580.
- [13] F. Riewe, *Nonconservative Lagrangian and Hamiltonian mechanics*, Phys. Rev. **E 53** (1996), 1890-1899.
- [14] F. Riewe, *Mechanics with fractional derivatives*. Phys. Rev. **E 55** (1997), 3581-3592.
- [15] E. Rabei, T. Alhalholy and A. Rousan, *Potential of arbitrary Forces with Fractional derivatives*, to appear in *Int. J. Theor. Phys. A*, (special communications).
- [16] E. Rabei, T. Alhalholy and A. Tanni, *On Hamiltonian formulation of nonconservative systems*, Turk. J. Phys. **28** (2004), 1-9.

- [17] O. P. Agrawal, *Formulation of Euler-Lagrange equations for fractional variational problems*, J. Math. Anal. Appl. **272** (2002), 368-379
- [18] O.P. Agrawal, *Lagrangian and Lagrange equation of motion for fractionally damped systems*, Trans. ASME, J. Math. Anal. Appl., **68** (2001) 339-341.
- [19] O. P. Agrawal, J. Gregory and K. P. Spector, *A bliss-type multiplier rule for constrained variational problems with time delay*, J. Math. Anal. Appl. **210**(1997) 702-711.
- [20] P. A. M. *Lectures on Quantum Mechanics* (Yeshiva University, New York, 1964); K. Sundermeyer, *Constrained Dynamics*, Lectures Notes in Physics Vol. 169 (Springer, New York, 1982) ; M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems* (Princeton Univ. Press, 1992); A. Hanson, T. Regge and C. Teitelboim, *Constrained Hamiltonian Systems*(Academia Nazionle dei Lincei, Rome, 1976)
- [21] D. Baleanu and T. Avkar, *Lagrangians with linear velocities within Riemann-Liouville fractional derivatives*, Nuovo cimento **119**, (2004) 73-79.
- [22] D. Baleanu, *Constrained systems and Riemann-Liouville fractional derivative*, Proceedings of 1st IFAC Workshop on Fractional Differentiation and its Applications, Bordeaux, France, July 19-21 (2004) 597-602.